

A NOTE ON POLY-BERNOULLI NUMBERS AND POLYNOMIALS OF THE SECOND KIND

TAEKYUN KIM, SANG HUN LEE, AND JONG JIN SEO

ABSTRACT. In this paper, we consider the poly-Bernoulli numbers and polynomials of the second kind and presents new and explicit formulae for calculating the poly-Bernoulli numbers of the second kind and the Stirling numbers of the second kind.

1. INTRODUCTION

As is well known, the Bernoulli polynomials of the second kind are defined by the generating function to be

$$(1) \quad \frac{t}{\log(1+t)}(1+t)^x = \sum_{n=0}^{\infty} b_n(x) \frac{t^n}{n!}, \quad (\text{see [5,14,16]}).$$

When $x = 0$, $b_n = b_n(0)$ are called the Bernoulli numbers of the second kind. The first few Bernoulli numbers b_n of the second kind are $b_0 = 1$, $b_1 = 1/2$, $b_2 = -1/12$, $b_3 = 1/24$, $b_4 = -19/720$, $b_5 = 3/160, \dots$

From (1), we have

$$(2) \quad b_n(x) = \sum_{l=0}^n \binom{n}{l} b_l (x)_{n-l},$$

where $(x)_n = x(x-1) \cdots (x-n+1)$, $(n \geq 0)$. The Stirling number of the second kind is defined by

$$(3) \quad x^n = \sum_{l=0}^n S_2(n, l)(x)_l, \quad (n \geq 0).$$

The ordinary Bernoulli polynomials are given by

$$(4) \quad \frac{t}{e^t - 1} e^{xt} = \sum_{n=0}^{\infty} B_n(x) \frac{t^n}{n!}, \quad (\text{see [1-18]}).$$

Key words and phrases. Bernoulli polynomials of the second, poly-Bernoulli numbers and polynomials, Stirling number of the second kind.

*Corresponding Author:

When $x = 0$, $B_n = B_n(0)$ are called the Bernoulli numbers.

It is known that the classical polylogarithmic function $Li_k(x)$ is given by

$$(5) \quad Li_k(x) = \sum_{n=1}^{\infty} \frac{x^n}{n^k}, (k \in \mathbb{Z}), \text{ (see [6,7,8])}.$$

For $k = 1$, $Li_1(x) = \sum_{n=1}^{\infty} \frac{x^n}{n} = -\log(1-x)$. The Stirling number of the first kind is defined by

$$(6) \quad (x)_n = \sum_{l=0}^n S_1(n, l) x^l, (n \geq 0), \text{ (see [15])}.$$

In this paper, we consider the poly-Bernoulli numbers and polynomials of the second kind and presents new and explicit formulae for calculating the poly-Bernoulli number and polynomial and the Stirling number of the second kind.

2. POLY-BERNOULLI NUMBERS AND POLYNOMIALS OF THE SECOND KIND

For $k \in \mathbb{Z}$, we consider the poly-Bernoulli polynomials $b_n^{(k)}(x)$ of the second kind as follows:

$$(7) \quad \frac{Li_k(1-e^{-t})}{\log(1+t)}(1+t)^x = \sum_{n=0}^{\infty} b_n^{(k)}(x) \frac{t^n}{n!}.$$

When $x = 0$, $b_n^{(k)} = b_n^{(k)}(0)$ are called the poly-Bernoulli numbers of the second kind.

Indeed, for $k = 1$, we have

$$(8) \quad \frac{Li_1(1-e^{-t})}{\log(1+t)}(1+t)^x = \frac{t}{\log(1+t)}(1+t)^x = \sum_{n=0}^{\infty} b_n(x) \frac{t^n}{n!}.$$

By (7) and (8), we get

$$(9) \quad b_n^{(1)}(x) = b_n(x), (n \geq 0).$$

It is known that

$$(10) \quad \frac{t(1+t)^{x-1}}{\log(1+t)} = \sum_{n=0}^{\infty} B_n^{(\alpha)}(x) \frac{t^n}{n!},$$

where $B_n^{(\alpha)}(x)$ are the Bernoulli polynomials of order α which are given by the generating function to be

$$\left(\frac{t}{e^t-1}\right)^{\alpha} e^{xt} = \sum_{n=0}^{\infty} B_n^{(\alpha)}(x) \frac{t^n}{n!}, \text{ (see [1-18])}.$$

By (1) and (10), we get

$$b_n(x) = B_n^{(n)}(x+1), (n \geq 0).$$

Now, we observe that

$$\begin{aligned}
 & \frac{Li_k(1 - e^{-t})}{\log(1 + t)} (1 + t)^x \\
 (11) \quad &= \sum_{n=0}^{\infty} b_n^{(k)}(x) \frac{t^n}{n!} \\
 &= \frac{1}{\log(1 + t)} \underbrace{\int_0^t \frac{1}{e^x - 1} \int_0^t \frac{1}{e^x - 1} \cdots \int_0^t \frac{1}{e^x - 1}}_{k-1 \text{ times}} \int_0^t \frac{x}{e^x - 1} dx \cdots dx (1 + t)^x.
 \end{aligned}$$

Thus, by (11), we get

$$\begin{aligned}
 \sum_{n=0}^{\infty} b_n^{(2)}(x) \frac{t^n}{n!} &= \frac{(1 + t)^x}{\log(1 + t)} \int_0^t \frac{x}{e^x - 1} dx \\
 (12) \quad &= \frac{(1 + t)^x}{\log(1 + t)} \sum_{l=0}^{\infty} \frac{B_l}{l!} \int_0^t x^l dx \\
 &= \left(\frac{t}{\log(1 + t)} \right) (1 + t)^x \sum_{l=0}^{\infty} \frac{B_l}{(l + 1) l!} t^l \\
 &= \sum_{n=0}^{\infty} \left\{ \sum_{l=0}^n \binom{n}{l} \frac{B_l b_{n-l}(x)}{l + 1} \right\} \frac{t^n}{n!}.
 \end{aligned}$$

Therefore, by (12), we obtain the following theorem.

Theorem 2.1. *For $n \geq 0$ we have*

$$b_n^{(2)}(x) = \sum_{l=0}^n \binom{n}{l} \frac{B_l b_{n-l}(x)}{l + 1}.$$

From (11), we have

$$\begin{aligned}
 \sum_{n=0}^{\infty} b_n^{(k)}(x) \frac{t^n}{n!} &= \frac{Li_k(1 - e^{-t})}{\log(1 + t)} (1 + t)^x \\
 (13) \quad &= \frac{t}{\log(1 + t)} \frac{Li_k(1 - e^{-t})}{t} (1 + t)^x.
 \end{aligned}$$

We observe that

$$\begin{aligned}
(14) \quad \frac{1}{t} Li_k(1 - e^{-t}) &= \frac{1}{t} \sum_{n=1}^{\infty} \frac{1}{n^k} (1 - e^{-t})^n \\
&= \frac{1}{t} \sum_{n=1}^{\infty} \frac{(-1)^n}{n^k} n! \sum_{l=n}^{\infty} S_2(l, n) \frac{(-t)^l}{l!} \\
&= \frac{1}{t} \sum_{l=1}^{\infty} \sum_{n=1}^l \frac{(-1)^{n+l}}{n^k} n! S_2(l, n) \frac{t^l}{l!} \\
&= \sum_{l=0}^{\infty} \sum_{n=1}^{l+1} \frac{(-1)^{n+l+1}}{n^k} n! \frac{S_2(l+1, n)}{l+1} \frac{t^l}{l!}.
\end{aligned}$$

Thus, by (10) and (14), we get

$$\begin{aligned}
(15) \quad \sum_{n=0}^{\infty} b_n^{(k)}(x) \frac{t^n}{n!} &= \left(\sum_{m=0}^{\infty} b_m(x) \frac{t^m}{m!} \right) \left\{ \sum_{l=0}^{\infty} \left(\sum_{p=1}^{l+1} \frac{(-1)^{p+l+1}}{p^k} p! \frac{S_2(l+1, p)}{l+1} \right) \frac{t^l}{l!} \right\} \\
&= \sum_{n=0}^{\infty} \left\{ \sum_{l=0}^n \binom{n}{l} \left(\sum_{p=1}^{l+1} \frac{(-1)^{p+l+1}}{p^k} p! \frac{S_2(l+1, p)}{l+1} \right) b_{n-l}(x) \right\} \frac{t^n}{n!}.
\end{aligned}$$

Therefore, by (15), we obtain the following theorem.

Theorem 2.2. *For $n \geq 0$, we have*

$$b_n^{(k)}(x) = \sum_{l=0}^n \binom{n}{l} \left(\sum_{p=1}^{l+1} \frac{(-1)^{p+l+1}}{p^k} p! \frac{S_2(l+1, p)}{l+1} \right) b_{n-l}(x).$$

By (7), we get

$$\begin{aligned}
(16) \quad \sum_{n=0}^{\infty} \left(b_n^{(k)}(x+1) - b_n^{(k)}(x) \right) \frac{t^n}{n!} &= \frac{Li_k(1 - e^{-t})}{\log(1+t)} (1+t)^{x+1} - \frac{Li_k(1 - e^{-t})}{\log(1+t)} (1+t)^x \\
&= \frac{t Li_k(1 - e^{-t})}{\log(1+t)} (1+t)^x \\
&= \left(\frac{t}{\log(1+t)} (1+t)^x \right) Li_k(1 - e^{-t}) \\
&= \left(\sum_{l=0}^{\infty} \frac{b_l(x)}{l!} t^l \right) \left\{ \sum_{p=1}^{\infty} \left(\sum_{m=1}^p \frac{(-1)^{m+p} m!}{m^k} S_2(p, m) \right) \right\} \frac{t^p}{p!}
\end{aligned}$$

$$\begin{aligned}
(17) \quad &= \sum_{n=1}^{\infty} \left(\sum_{p=1}^n \sum_{m=1}^p \frac{(-1)^{m+p}}{m^k} m! S_2(p, m) \frac{b_{n-p}(x) n!}{(n-p)! p!} \right) \frac{t^n}{n!} \\
&= \sum_{n=1}^{\infty} \left\{ \sum_{p=1}^n \sum_{m=1}^p \binom{n}{p} \frac{(-1)^{m+p} m!}{m^k} S_2(p, m) b_{n-p}(x) \right\} \frac{t^n}{n!}.
\end{aligned}$$

Therefore, by (16), we obtain the following theorem.

Theorem 2.3. *For $n \geq 1$, we have*

$$(18) \quad b_n^{(k)}(x+1) - b_n^{(k)}(x) = \sum_{p=1}^n \sum_{m=1}^p \binom{n}{p} \frac{(-1)^{m+p} m!}{m^k} S_2(p, m) b_{n-p}(x).$$

From (13), we have

$$\begin{aligned}
(19) \quad &\sum_{n=0}^{\infty} b_n^{(k)}(x+y) \frac{t^n}{n!} = \left(\frac{Li_k(1-e^{-t})}{\log(1+t)} \right)^k (1+t)^{x+y} \\
&= \left(\frac{Li_k(1-e^{-t})}{\log(1+t)} \right)^k (1+t)^x (1+t)^y \\
&= \left(\sum_{l=0}^{\infty} b_l^{(k)}(x) \frac{t^l}{l!} \right) \left(\sum_{m=0}^{\infty} (y)_m \frac{t^m}{m!} \right) \\
&= \sum_{n=0}^{\infty} \left(\sum_{l=0}^n (y)_l b_{n-l}^{(k)}(x) \frac{n!}{(n-l)! l!} \right) \frac{t^n}{n!} \\
&= \sum_{n=0}^{\infty} \left(\sum_{l=0}^n \binom{n}{l} b_{n-l}^{(k)}(x) (y)_l \right) \frac{t^n}{n!}.
\end{aligned}$$

Therefore, by (17), we obtain the following theorem.

Theorem 2.4. *For $n \geq 0$, we have*

$$b_n^{(k)}(x+y) = \sum_{l=0}^n \binom{n}{l} b_{n-l}^{(k)}(x) (y)_l.$$

Acknowledgements. The present Research has been conducted by the Research Grant of Kwangwoon University in 2014

REFERENCES

- [1] J. Choi, D. S. Kim, T. Kim, and Y. H. Kim, Some arithmetic identities on Bernoulli and Euler numbers arising from the p -adic integrals on Z_p , *Adv. Stud. Contemp. Math.* **22** (2012), 239-247.
- [2] D. Ding and J. Yang, Some identities related to the Apostol-Euler and Apostol-Bernoulli polynomials, *Adv. Stud. Contemp. Math.* **20** (2010), 7-21.
- [3] S. Gaboury, R. Tremblay, and B.-J. Fugère, Some explicit formulas for certain new classes of Bernoulli, Euler and Genocchi polynomials, *Proc. Jangjeon Math. Soc.* **17** (2014), 115-123.
- [4] D. King, S. J. Lee, L.-W. Park, and S.-H. Rim, On the twisted weak weight q -Bernoulli polynomials and numbers, *Proc. Jangjeon Math. Soc.* **16** (2013), 195-201.
- [5] D. S. Kim, T. Kim, and S.-H. Lee, Poly-Cauchy numbers and polynomials with umbral calculus viewpoint, *Int. J. Math. Anal.* **7** (2013), 2235-2253.
- [6] D. S. Kim, T. Kim, and S.-H. Lee, A note on poly-Bernoulli polynomials arising from umbral calculus, *Adv. Stud. Theor. Phys.* **7** (2013), no. 15, 731-744.
- [7] D. S. Kim and T. Kim, Higher-order Frobenius-Euler and poly-Bernoulli mixed-type polynomials, *Adv. Difference Equ.* **2013**, 2013:251, 13 pp.
- [8] D. S. Kim, T. Kim, S.-H. Lee, and S.-H. Rim, Umbral calculus and Euler polynomials, *Ars Combin.* **112** (2013), 293-306.
- [9] D. S. Kim and T. Kim, Higher-order Cauchy of first kind and poly-Cauchy of the first kind mixed type polynomials, *Adv. Stud. Contemp. Math.* **23** (2013), no. 4, 621-636.
- [10] T. Kim, q -Bernoulli numbers and polynomials associated with Gaussian binomial coefficients, *Russ. J. Math. Phys.* **15** (2008), 51-57.
- [11] Y.-H. Kim and K.-W. Hwang, Symmetry of power sum and twisted Bernoulli polynomials, *Adv. Stud. Contemp. Math.* **18** (2009), 127-133.
- [12] H. Ozden, I. N. Cangul, and Y. Simsek, Remarks on q -Bernoulli numbers associated with Daehee numbers, *Adv. Stud. Contemp. Math.* **18** (2009), 41-48.
- [13] J.-W. Park, New approach to q -Bernoulli polynomials with weight or weak weight, *Adv. Stud. Contemp. Math.* **24** (2014), no. 1, 39-44.
- [14] T. R. Prabhakar and S. Gupta, Bernoulli polynomials of the second kind and general order, *Indian J. Pure Appl. Math.* **11** (1980), 1361-1368.
- [15] S. Roman, *The Umbral Calculus. Pure and Applied Mathematics*, Vol. 111, Academic Press, New York (1984), ISBN:0-12-594380-6.
- [16] S. Roman and G. C. Rota, The umbral calculus, *Adv. Math.* **27** (1978), no. 2, 95-188.
- [17] Y. Simsek, Generating functions of the twisted Bernoulli numbers and polynomials associated with their interpolation functions, *Adv. Stud. Contemp. Math.* **16** (2008), 251-278.
- [18] H. M. Srivastava, T. Kim, and Y. Simsek, q -Bernoulli numbers and polynomials associated with multiple q -zeta functions and basic L -series, *Russ. J. Math. Phys.* **12** (2005), 241-278.

T. KIM
 DEPARTMENT OF MATHEMATICS
 KWANGWOON UNIVERSITY
 SEOUL 139-701, REPUBLIC OF KOREA
 e-mail: tkkim@kw.ac.kr

S. H. LEE
 DIVISION OF GENERAL EDUCATION
 KWANGWOON UNIVERSITY
 SEOUL 139-701, REPUBLIC OF KOREA
 e-mail: tkkim@kw.ac.kr

J. J. SEO
DEPARTMENT OF APPLIED MATHEMATICS
PUKYONG NATIONAL UNIVERSITY
PUSAN 698-737, REPUBLIC OF KOREA
e-mail: seo2011@pknu.ac.kr